

STAT 309: MATHEMATICAL COMPUTATIONS I
FALL 2022
PROBLEM SET 0

This homework mostly serves as a linear algebra refresher. We will recall some definitions. The null space or kernel of a matrix $A \in \mathbb{R}^{m \times n}$ is the set

$$\ker(A) = \{\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} = \mathbf{0}\}$$

while the range space or image is the set

$$\operatorname{im}(A) = \{\mathbf{y} \in \mathbb{R}^m : \mathbf{y} = A\mathbf{x} \text{ for some } \mathbf{x} \in \mathbb{R}^n\}.$$

The rank and nullity of A are defined as the dimensions of these spaces,

$$\operatorname{rank}(A) = \dim \operatorname{im}(A) \quad \text{and} \quad \operatorname{nullity}(A) = \dim \ker(A).$$

By convention we write all vectors in \mathbb{R}^n as column vectors.

1. (a) For $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{n \times p}$, show that

$$\operatorname{im}(AB) \subseteq \operatorname{im}(A) \quad \text{and} \quad \ker(AB) \supseteq \ker(B).$$

When does equality occur in each of these inclusions?

- (b) For $A, B \in \mathbb{R}^{n \times n}$, show that

$$\begin{aligned} \operatorname{rank}(AB) &\leq \min\{\operatorname{rank}(A), \operatorname{rank}(B)\}, \\ \operatorname{nullity}(AB) &\leq \operatorname{nullity}(A) + \operatorname{nullity}(B), \\ \operatorname{rank}(A + B) &\leq \operatorname{rank}(A) + \operatorname{rank}(B). \end{aligned}$$

- (c) For $A, B \in \mathbb{R}^{n \times n}$, show that if $AB = 0$, then

$$\operatorname{rank}(A) + \operatorname{rank}(B) \leq n.$$

2. (a) Let $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{p \times q}$. Show that

$$\operatorname{rank} \left(\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \right) = \operatorname{rank}(A) + \operatorname{rank}(B).$$

We have used the block matrix notation here. For example if $A = \begin{bmatrix} a & b & c \\ d & e & f \end{bmatrix} \in \mathbb{R}^{2 \times 3}$ and $B = \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \in \mathbb{R}^{2 \times 1}$, then

$$\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} = \begin{bmatrix} a & b & c & 0 \\ d & e & f & 0 \\ 0 & 0 & 0 & \alpha \\ 0 & 0 & 0 & \beta \end{bmatrix} \in \mathbb{R}^{4 \times 4}.$$

This is sometimes also denoted as $A \oplus B$. It is a direct sum of operators induced by a direct sum of vector spaces.

- (b) For $\mathbf{x} = [x_1, \dots, x_m]^T \in \mathbb{R}^m$ and $\mathbf{y} = [y_1, \dots, y_n]^T \in \mathbb{R}^n$, observe that $\mathbf{xy}^T \in \mathbb{R}^{m \times n}$. Let $A \in \mathbb{R}^{m \times n}$. Show that $\operatorname{rank}(A) = 1$ iff $A = \mathbf{xy}^T$ for some nonzero $\mathbf{x} \in \mathbb{R}^m$ and $\mathbf{y} \in \mathbb{R}^n$.

3. Let $A \in \mathbb{R}^{m \times n}$.

(a) Show that

$$\ker(A^T A) = \ker(A) \quad \text{and} \quad \text{im}(A^T A) = \text{im}(A^T).$$

Give an example to show this is not true over a finite field (e.g. a field of two elements $\mathbb{F}_2 = \{0, 1\}$ with binary arithmetic).

(b) Show that

$$A^T A \mathbf{x} = A^T \mathbf{b}$$

always has a solution (even if $A \mathbf{x} = \mathbf{b}$ has no solution). Give an example to show that this is not true over a finite field.

4. Let $\mathbf{v}_1, \dots, \mathbf{v}_n \in \mathbb{R}^n$. Let $r \leq n$ and $G_r = [g_{ij}] \in \mathbb{R}^{r \times r}$ be the matrix with

$$g_{ij} = \mathbf{v}_i^T \mathbf{v}_j$$

for $i, j = 1, \dots, r$. This is called a *Gram matrix*.

(a) Show that $\mathbf{v}_1, \dots, \mathbf{v}_r$ are linearly independent iff $\text{nullity}(G_r) = 0$.

(b) Show that $G_r = I_r$ iff $\mathbf{v}_1, \dots, \mathbf{v}_r$ are pairwise orthogonal unit vectors, i.e., $\|\mathbf{v}_i\|_2 = 1$ for all $i = 1, \dots, r$, and $\mathbf{v}_i^T \mathbf{v}_j = 0$ for all $i \neq j$. If this holds, show that

$$\sum_{i=1}^r (\mathbf{v}^T \mathbf{v}_i)^2 \leq \|\mathbf{v}\|_2^2 \quad (4.1)$$

for all $\mathbf{v} \in \mathbb{R}^n$. What can you say about $\mathbf{v}_1, \dots, \mathbf{v}_r$ if equality always holds in (4.1) for all $\mathbf{v} \in \mathbb{R}^n$?

5. Let $A, B \in \mathbb{R}^{m \times n}$. For any $i, j \in \{1, \dots, n\}$, let $E_{ij} \in \mathbb{R}^{p \times q}$ denote the matrix with one in the (i, j) th entry and zeros everywhere else.

(a) Describe the matrix $E_{ij} A E_{kl} \in \mathbb{R}^{m \times n}$ where $E_{ij} \in \mathbb{R}^{p \times m}$ and $E_{kl} \in \mathbb{R}^{n \times q}$.

(b) Find expressions $\text{tr}(A^T B)$, $\text{tr}(AB^T)$, $\text{tr}(BA^T)$, $\text{tr}(B^T A)$ in terms of the entries of A and B . What if $A = B$? What if $B = E_{ij} \in \mathbb{R}^{m \times n}$?

(c) Suppose $m = n$ and $\text{tr}(A) = \text{tr}(A^2) = \dots = \text{tr}(A^n) = 0$. Show that $A^n = 0$.

(d) Suppose $m = n$. Show that (i) $\text{tr}[(AB)^k] = \text{tr}[(BA)^k]$ and (ii) if $AB = 0$, then $\text{tr}[(A+B)^k] = \text{tr}(A^k) + \text{tr}(B^k)$, for all $k \in \mathbb{N}$.

6. Let $A \in \mathbb{C}^{n \times n}$. Recall that A is diagonalizable iff there exists an invertible $X \in \mathbb{C}^{n \times n}$ such that $X^{-1} A X = \Lambda$, a diagonal matrix.

(a) Show that A is diagonalizable if and only if its minimal polynomial is of the form

$$m_A(x) = (x - \lambda_1) \cdots (x - \lambda_d)$$

where $\lambda_1, \dots, \lambda_d \in \mathbb{C}$ are all distinct. Hence deduce for a diagonalizable matrix, the degree of its minimal polynomial equals the number of distinct eigenvalues.

(b) Let A be diagonalizable. Let $\mathbf{x}_1, \dots, \mathbf{x}_n \in \mathbb{C}^n$ be n linearly independent right eigenvectors, i.e., $A \mathbf{x}_i = \lambda_i \mathbf{x}_i$; and $\mathbf{y}_1, \dots, \mathbf{y}_n \in \mathbb{C}^n$ be n linearly independent left eigenvectors, i.e., $\mathbf{y}_i^T A = \lambda_i \mathbf{y}_i^T$. Show that we may choose $\mathbf{x}_1, \dots, \mathbf{x}_n$ and $\mathbf{y}_1, \dots, \mathbf{y}_n$ so that any vector $\mathbf{v} \in \mathbb{C}^n$ can be expressed as

$$\mathbf{v} = \sum_{i=1}^n (\mathbf{y}_i^T \mathbf{v}) \mathbf{x}_i.$$

If we write $X = [\mathbf{x}_1, \dots, \mathbf{x}_n] \in \mathbb{C}^{n \times n}$ and $Y = [\mathbf{y}_1, \dots, \mathbf{y}_n] \in \mathbb{C}^{n \times n}$. What is the relation between X and Y ?